## **Amendments to the Claims:**

This listing of claims will replace all prior versions, and listings, of claims in the application:

## **Listing of Claims:**

Claims 1-7 (canceled)

- 8. (withdrawn) A DNA fragment having an initiation codon, a stop codon and a coding sequence between said two codons, said coding sequence substantially corresponding to said amino acid sequence of claim 1.
- 9. (withdrawn) The DNA fragment of claim 8, wherein said DNA fragment has a sequence of nucleotide residues substantially identical to SEQ ID NO: 4 in FIG 2.
- 10. (withdrawn) The DNA fragment of claim 8, wherein said DNA fragment has a sequence of nucleotide residues substantially identical to SEQ ID NO: 5 in FIG 3.
- 11. (withdrawn) A method of producing said truncated glucanase of claim 1, comprising:
- (a) growing in a culture medium a bacterial strain containing a gene encoding for a wild-type 1,3-1,4-β-D-glucanase from *Fibrobacter succinogenes*,
  - (b) centrifuging said culture medium to produce a supernatant,
  - (c) incubating said supernatant to produce said truncated glucanase, and
  - (d) collecting and purifying said truncated glucanase from said supernatant.

- 12. (withdrawn) The method of claim 11, wherein said supernatant in step (d) is incubated for at least 7 days at 4 °C or a higher temperature.
- 13. (withdrawn) The method of claim 11, wherein said supernatant in step (d) is incubated for a period ranging from 10 days to 14 days and at a temperature ranging from 4 °C to 37 °C.
- 14. (withdrawn) The method of claim 11, wherein said supernatant in step (d) is incubated for 14 days at 37 °C.
- 15. (withdrawn) A method of producing said truncated glucanase of claim 1, comprising:
- (a) amplifying a DNA fragment using a PCR method from a DNA template containing a gene encoding for a wild-type glucanase from *Fibrobacter sucinogenes*, said DNA fragment substantially corresponding to a portion of said gene,
  - (b) subcloning said amplified DNA fragment in an expression vector,
- (c) transferring said expression vector harbouring said DNA fragment into a host strain,
- (d) growing said host strain in a culture medium for a period of time and inducing expression of said DNA fragment, with or without adding an inducer, to produce a sufficient amount of protein products, and
  - (e) collecting and purifying protein expression products from said culture medium.

- 16. (withdrawn) The method of claim 15, wherein said DNA fragment amplified in step (a) has a sequence substantially identical to SEQ ID NO: 6 in FIG. 6.
- 17. (withdrawn) The method of claim 11, wherein said gene encoding for a wild-type 1,3-1,4-β-D-glucanase is carried in a plasmid.
- 18. (withdrawn) The method of claim 17, further comprising, between step(a) and step(b), an additional step of adding to said culture medium an inducer to induce expression of said gene.
  - 19. (withdrawn) The method of claim 15, wherein said host strain is a bacterial strain.
- 20. (currently amended) An isolated truncated glucanase having enhanced glucanase activity relative to a matured wild type glucanase and an amino acid sequence of a total number of amino acid residues between 248 to 322 less than 322 amino acid residues, comprising a portion of SEQ ID NO:3, said portion of SEQ ID NO: 3 beginning at amino acid residue position 28th and ending at amino acid residue position 271th of said SEQ ID NO: 3. wherein a portion of said amino acid sequence starting at amino acid residue 5th and extending at least 243 amino acid therefrom residue is substantially identical to a corresponding portion of SEQ ID No: 3, starting at amino acid residue 28th of SEQ ID No. 3 and extending therefrom.
- 21. (previously presented) The isolated truncated glucanase of claim 20, absent a repeated PXSSSS segment, wherein X represents an uncharged amino acid residue.

- 22. (currently amended) The SEQ ID No. 3 of claim 20 The isolated glucanase of claim 20, wherein the SEQ ID No: 3 is identical to a wild-type glucanase from Fibrobacter Fibrobactor succinogenes.
- 23. (previously presented) The isolated truncated glucanase of claim 20 having an amino acid sequence substantially identical to SEQ ID No: 1.
- 24. (previously presented) The isolated truncated glucanase of claim 20 having an amino acid sequence substantially identical to SEQ ID No: 2.